

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Identification of biomarkers of colitis to monitor effects of dietary
omega-3 polyunsaturated fatty acids in the interleukin-10 gene-
deficient mouse model of inflammatory bowel diseases**

A thesis presented in partial fulfilment of the requirements for the

degree of

Doctor of Philosophy

in

Nutritional Science

at Massey University, Manawatū,

New Zealand.

Nadja Berger

2016

ABSTRACT

Inflammatory bowel diseases (IBD) are characterised by chronic inflammation of the gastrointestinal tract including the colon (colitis). Increased dietary intake of salmon, which is rich in eicosapentaenoic acid (EPA), was well tolerated by IBD patients, leading to a perceived decrease in symptoms. However, better knowledge of the mechanisms by which EPA-rich diets affect IBD severity, and appropriate biomarkers for assessing these effects, are needed for potential targeted nutritional interventions.

This dissertation aimed to determine the temporal effects (early (9 weeks of age) vs. established (12 weeks)) of a diet containing 3.7% EPA, and the dose-dependent effects (15% to 45%) of a salmon diet at 12 weeks of age, on the severity of colitis. Molecular responses in colon and/or liver of the interleukin-10 gene-deficient (*Il10^{-/-}*) mouse model of IBD and healthy mice were assessed. Caecum digesta, urine and blood were mined to identify biomarkers (microbiota, metabolites and genes) of these responses.

The EPA diet reduced the severity of colitis only in 12-week-old *Il10^{-/-}* mice. This response was associated with changes in gene expression associated with lymphocyte function, eicosanoid signalling and peroxisome proliferator-activated receptor gamma signalling. The blood immune cell gene expression profile did not correlate with reduced colitis in these mice, but the urine metabolite profile was related to changes in colonic tryptophan metabolism.

The effects of the salmon diets on colitis were dose-dependent in 12-week-old *Il10^{-/-}* mice. The intermediate amount of salmon (30%) reduced the severity of colitis and lymphocyte-related gene expression, while enhancing genes in metabolic pathways. Tryptophan metabolism was not affected in these mice, but the urinary metabolite profile correlated with effects on hepatic tocopherol metabolism, as shown by reduced abundance of gamma-carboxyethylhydroxychroman glucoside. The abundances of *V. akkermansia*, *Eubacterium* spp., and an unclassified *Rikenellaceae* were further affected in these mice.

This is the first report describing molecular responses in the colon and liver of *Il10^{-/-}* mice fed a salmon diet associated with reduced colitis. Ultimately these responses could be validated for use in humans, and potentially enable management of IBD with diet.

ACKNOWLEDGEMENTS

I would like to express my sincerest gratitude to my supervisors Assoc Prof Nicole Roy, Dr Emma Bermingham, Prof Warren McNabb (all AgResearch Grasslands, Palmerston North, NZ) and Dr Janine Cooney (Plant & Food Research Ruakura, Hamilton, NZ) for your guidance and support throughout this project. Thank you for your encouragement, wisdom and finally your patience. I could not possibly have reached this point without you. I also acknowledge AgResearch for funding this project within the Nutrigenomics New Zealand partnership.

I would like to thank Dr Matthew Barnett (AgResearch Grasslands) for letting me be part of the “EPA time-course experiment” after one of my own mouse experiments had failed. Your enthusiasm and positivity was much appreciated when I did not know how to continue.

I am grateful to Dr Wayne Young (AgResearch Grasslands), an expert in ‘Omics’ and microbiota analyses, for providing various codes in R and helping with all enquiries. Also, thank you for proof-reading parts of this dissertation and providing valuable feedback.

Proteomics work was performed at Plant & Food Research in Auckland under the guidance of Ms Di Brewster. Thank you Di for taking the time to teach the art of 2D-DIGE and your ongoing technical advice. Also, thanks to all members from this laboratory for your welcoming attitudes which made an otherwise difficult process great. I am also grateful to Dr Janine Cooney for conducting MS identification.

A number of people behind this project deserve to be both acknowledged and thanked here: Mr Shuotun Zhu (University of Auckland, Auckland, NZ) for histological analysis. Ms Kelly Armstrong for the training in histology, RNA extraction and microarray analysis. Dr John Koolaard (AgResearch Grasslands) for statistical advice. Mr Paul Maclean (AgResearch Ruakura) for assistance with microarray data processing. Mr Jason Peters and Ms Leigh Ryan for help with animal work (AgResearch Grasslands). Mr Bruce Sinclair (AgResearch Grasslands) for reading various drafts of this dissertation and lending a helping hand during experiments. Dr Mark McCann (AgResearch Grasslands) for assistance with qPCR validation. Metabolomics experts Dr Karl Fraser and Dr Jan Huege (who sadly passed away during the course of this project) for their

valuable advice and sharing their expertise (AgResearch Grasslands). Thank you all for your help and contribution!

I am grateful to Prof Chris Evelo for giving me the opportunity to conduct parts of my last research chapter in the BigCaT Bioinformatics Group at the University of Maastricht in the Netherlands and would like to thank all the members from this group for making my stay in Maastricht a great one. Special thanks to Dr Lars Eijssen for assistance with data analysis, sharing his knowledge in R and the good times biking through the streets of Maastricht on the search for a beer (or two).

I have been very lucky to go through this process with a group of people that provided moral support and most importantly their friendship. Thank you to Ms Denise Martin (AgResearch Grasslands) for your encouraging words when I had already given up. You are an incredible person and have become a true friend. A huge thanks to my fellow postgraduate students for the chats around the lunch table and the many laughs we shared during road trips to Auckland or a girls-night-out. You truly made the day more enjoyable.

Finally I would like to thank my family with all my heart. My parents Nora and Wolfgang for your ongoing love and support. My brother Gerold for watching over me since 1987. My granny Waldtraud and aunt Traudi for your generosity. To my partner Paul thank you for being absolutely amazing throughout the last few years.

TABLE OF CONTENTS

Abstract	i
Acknowledgements	iii
Table of Contents	v
List of Figures	xi
List of Tables	xiv
List of Abbreviations	xvii
List of Appendices	xvi
Introduction	xix
Review of literature	1
1.1 Inflammatory bowel diseases	2
1.1.1 Genetic susceptibility	4
1.1.2 Mucosal immune-regulation	4
1.1.2.1 Intestinal epithelium	5
1.1.2.2 Intestinal inflammation	8
1.1.2.3 The commensal microbiota	10
1.1.3 The role of the environment	14
1.1.4 Animal models of IBD	15
1.1.4.1 The interleukin-10 gene-deficient mouse model	18
1.2 The concept of systems biology	19
1.2.1 Nutrigenomics	21
1.2.2 Exploration of biomarkers of intestinal inflammation	21
1.3 Dietary salmon and intestinal inflammation	23
1.4 Anti-inflammatory effects of specific salmon components	24
1.4.1 Micronutrients	24
1.4.2 Peptides	26
1.4.3 Lipids	26
1.4.4 Putative mechanisms of action of lipids	31
1.4.4.1 Modulation of cell membrane lipid rafts	31
1.4.4.2 Formation of lipid mediators	33
1.4.4.3 Modulation of gene expression	33
1.4.4.4 Modulation of protein expression	35
1.4.4.5 Modulation of the intestinal microbiota	36
1.5 Conclusion and outlook	37
1.6 Hypothesis and aims of the dissertation	38
1.7 Approach and structure of the dissertation	38
Materials and Methods	41
2.1 Introduction	42
2.2 EPA time-course experiment	43
2.2.1 Experimental design	43
2.2.2 Mouse model and induction of colitis	43
2.2.3 Experimental diets	45
2.2.4 Sampling procedure and tissue collection	46
2.2.5 Statistical evaluation of growth performance	48

2.3	Salmon diet experiment	48
2.3.1	Experimental design	48
2.3.2	Mouse model and induction of colitis	48
2.3.3	Experimental diets	50
2.3.4	Sampling procedure and tissue collection	51
2.3.5	Statistical evaluation of growth performance	53
2.4	Histopathological assessment	53
2.4.1	Tissue preparation and haematoxylin and eosin stain	53
2.4.2	Histological injury score	53
2.4.3	Statistical evaluation of histopathological changes	54
2.5	Isolation of peripheral blood mononuclear cells and RNA extraction	55
2.6	Extraction of RNA and protein from colon and liver	56
2.7	Transcriptomic analysis of colon, liver and peripheral blood mononuclear cells	57
2.7.1	Method overview	57
2.7.2	Sample preparation	57
2.7.3	CyDye-labelling and microarray hybridisation	59
2.7.4	Microarray data processing	60
2.8	Proteomic analysis of colon tissue	61
2.8.1	Method overview	62
2.8.2	Sample preparation	65
2.8.3	CyDye-labelling	67
2.8.4	Separation in the first dimension	67
2.8.5	Equilibration	68
2.8.6	Separation in the second dimension	68
2.8.7	In-gel protein digestion	68
2.8.8	Protein identification	69
2.9	Metabolomic analysis of urine	70
2.9.1	Method overview	71
2.9.2	Sample preparation and LC-MS analysis	71
2.9.3	MS data processing and statistical analysis	73
2.10	Microbiomic analysis of caecum digesta	77
2.10.1	Method overview	77
2.10.2	Extraction of metagenomic DNA and 454 pyrosequencing	79
2.10.3	Data analysis	79
2.11	Bioinformatic analysis	80
2.11.1	Ingenuity Pathway Analysis	80
2.11.2	Gene Set Enrichment Analysis	81
2.11.3	Integration of 'Omics' data	82
2.12	Conclusion and outlook	82

Effects of eicosapentaenoic acid-based diets on early and established colitis in the interleukin-10 gene-deficient mouse **87**

3.1	Introduction	88
3.2	Hypothesis and aim	89
3.3	Methods	90
3.4	Results	91
3.4.1	Growth performance	91
3.4.2	Severity of intestinal inflammation at 9 and 12 weeks of age	95
3.4.3	Colon gene expression	98

3.4.3.1	Comparison of colon gene expression between the AIN-76A and OA diet in <i>Il10</i> ^{-/-} mice	98
3.4.3.2	Colon gene expression between mouse genotypes fed the OA diet	98
3.4.3.3	Effect of the EPA diet (vs. OA diet) on colon gene expression in <i>Il10</i> ^{-/-} mice	105
3.4.4	PBMC gene expression	115
3.4.4.1	PBMC gene expression between mouse genotypes fed the OA diet	118
3.4.4.2	Effect of the EPA diet (vs. OA diet) on PBMC gene expression in <i>Il10</i> ^{-/-} mice	119
3.4.5	Comparison of colon and PBMC gene expression	130
3.4.5.1	Colon and PBMC gene expression between mouse genotypes fed the OA diet	130
3.4.5.2	Effect of the EPA diet (vs. OA diet) on colon and PBMC gene expression in <i>Il10</i> ^{-/-} mice	136
3.4.6	Urinary metabolites	136
3.4.6.1	Urinary metabolites between mouse genotypes irrespective of diet	139
3.4.6.2	Effect of the EPA diet (vs. OA diet) on urinary metabolites irrespective of genotype	150
3.4.6.3	Effect of the EPA diet (vs. OA diet) on urinary metabolites in <i>Il10</i> ^{-/-} mice	157
3.5	Discussion	161
3.5.1	Growth performance	162
3.5.2	Severity of colitis in <i>Il10</i> ^{-/-} mice	163
3.5.3	Transcriptomic profiling of PBMCs and colon tissue	163
3.5.3.1	Colon transcriptomic profile during colitis	163
3.5.3.2	Colon transcriptomic profile in response to the EPA diet (vs. OA diet) in <i>Il10</i> ^{-/-} mice	166
3.5.3.3	PBMC transcriptomic profile during colitis	169
3.5.3.4	PBMC transcriptomic profile in response to the EPA diet (vs. OA diet) in <i>Il10</i> ^{-/-} mice	170
3.5.4	Urine metabolomics	171
3.5.4.1	Urine metabolomic profile during colitis	171
3.5.4.2	Urine metabolomic profile in response to the EPA diet (vs. OA diet) irrespective of genotype	172
3.5.4.3	Urine metabolomic profile in response to the EPA diet (vs. OA diet) in <i>Il10</i> ^{-/-} mice	173
3.6	Conclusion and outlook	174

Dose-response of salmon-based diets on established colitis and associated colonic gene expression in the interleukin-10 gene-deficient mouse **175**

4.1	Introduction	176
4.2	Hypothesis and aim	177
4.3	Methods	177
4.4	Results	179
4.4.1	Experimental diet composition	179
4.4.2	Growth performance	181
4.4.3	Severity of intestinal inflammation	184
4.4.4	Colon gene expression	187
4.4.4.1	Colon gene expression between mouse genotypes fed the AIN-76A diet	187
4.4.4.2	Colon gene expression in mice fed the 15% salmon and 15% control diets	193
4.4.4.2.1	Colon gene expression between mouse genotypes fed the 15% control diet	193

4.4.4.2.2	Effect of the 15% salmon diet (vs. 15% control diet) on colon gene expression in <i>Il10</i> ^{-/-} mice	197
4.4.4.3	Colon gene expression in mice fed the 30% salmon and 30% control diets	200
4.4.4.3.1	Colon gene expression between mouse genotypes fed the 30% control diet	200
4.4.4.3.2	Effect of the 30% salmon diet (vs. 30% control diet) on colon gene expression in <i>Il10</i> ^{-/-} mice	205
4.4.4.4	Colon gene expression in mice fed the 45% salmon and 45% control diets	207
4.4.4.4.1	Colon gene expression between mouse genotypes fed the 45% control diet	212
4.4.4.4.2	Effect of the 45% salmon diet (vs. 45% control diet) on colon gene expression irrespective of genotype	214
4.5	Discussion	220
4.5.1	Levels of LC n-3 PUFA	221
4.5.2	Severity of colitis in <i>Il10</i> ^{-/-} mice	222
4.5.3	Transcriptomic profiling of colon tissue	224
4.5.3.1	Pro-inflammatory gene expression in <i>Il10</i> ^{-/-} mice fed the 15% salmon diet (vs. 15% control diet)	224
4.5.3.2	Enhanced metabolic pathways in <i>Il10</i> ^{-/-} mice fed the 30% salmon diet (vs. 30% control diet)	225
4.5.3.3	Effect of the 45% salmon diet (vs. 45% control diet) dependent on genotype	228
4.6	Conclusion and outlook	229

Multi-‘Omics’ approach to investigate the effects of a diet containing 30% salmon on the microbial community in the caecum and immune and metabolic pathways in colon and liver in the interleukin-10 gene-deficient mouse **231**

5.1	Introduction	232
5.2	Hypothesis and aim	233
5.3	Methods	234
5.4	Results	234
5.4.1	Colon protein expression	234
5.4.1.1	Identification of proteins	234
5.4.1.2	Colon protein expression between mouse genotypes fed the 30% control diet	241
5.4.1.3	Effect of the 30% salmon diet (vs. 30% control diet) on colon protein expression in <i>Il10</i> ^{-/-} mice	243
5.4.2	Liver gene expression	244
5.4.3.1	Liver gene expression between mouse genotypes fed the 30% control diet	247
5.4.3.2	Effect of the 30% salmon diet (vs. 30% control diet) on liver gene expression in <i>Il10</i> ^{-/-} mice	252
5.4.4	Urinary metabolites	255
5.4.4.1	Metabolite identification	255
5.4.4.2	Metabolomic fingerprinting	259
5.4.4.2.1	Urinary metabolites between mouse genotypes fed the 30% control diet	263
5.4.4.2.2	Effect of the 30% salmon diet (vs. 30% control diet) on urinary metabolites in <i>Il10</i> ^{-/-} mice	266
5.4.5	Analysis of microbiota from caecum digesta	267
5.4.5.1	Species diversity estimate	267

5.4.5.2	Taxonomic differences	267
5.4.5.3	Beta diversity	273
5.5	Discussion	273
5.5.1	Proteomic profiling of colon tissue	276
5.5.1.1	Colon proteomic profile during colitis	276
5.5.1.2	Colon proteomic profile in response to the 30% salmon diet (vs. 30% control diet) in <i>Il10</i> ^{-/-} mice	277
5.5.2	Transcriptomic profiling of liver tissue	280
5.5.2.1	Liver transcriptomic profile during colitis	280
5.5.2.2	Liver transcriptomic profile in response to the 30% salmon diet (vs. 30% control diet) irrespective of genotype	281
5.5.2.3	Liver transcriptomic profile in response to the 30% salmon diet (vs. 30% control diet) in <i>Il10</i> ^{-/-} mice	283
5.5.3	Urine metabolomics	284
5.5.3.1	Urine metabolomic profile during colitis	284
5.5.3.2	Urine metabolomic profile in response to the 30% salmon diet (vs. 30% control diet) irrespective of genotype	285
5.5.3.3	Urine metabolomic profile in response to the 30% salmon diet (vs. 30% control diet) in <i>Il10</i> ^{-/-} mice	286
5.5.4	Microbiomic analysis of caecum digesta	287
5.5.4.1	Microbial community profile during colitis	287
5.5.4.2	Microbial community profile in response to the 30% salmon diet (vs. 30% control diet) irrespective of genotype	287
5.5.4.3	Microbial community profile in response to the 30% salmon diet (vs. 30% control diet) in <i>Il10</i> ^{-/-} mice	288
5.6	Conclusion and outlook	289
Integration of ‘Omics’ data to characterise the systemic responses to a diet containing 30% salmon in the interleukin-10 gene-deficient mouse		291
6.1	Introduction	292
6.2	Hypothesis and aim	293
6.3	Methods	293
6.4	Results	294
6.4.1	Tryptophan metabolism between mouse genotypes fed the 30% control diet	294
6.4.2	Tocopherol metabolism between mouse genotypes fed the 30% control diet	294
6.4.3	Microbial community profile between mouse genotypes fed the 30% control diet	295
6.4.4	Colon and hepatic gene expression and microbial community profile and their relationship to the urinary metabolite profile in <i>Il10</i> ^{-/-} mice fed the 30% salmon diet (vs. 30% control diet)	300
6.5	Discussion	305
6.5.1	Novel insights into tryptophan metabolism during colitis	305
6.5.2	Novel insights into tocopherol metabolism during colitis	306
6.5.3	Novel insights into tocopherol metabolism in response to the 30% salmon diet (vs. 30% control diet) in <i>Il10</i> ^{-/-} mice	307
6.6	Conclusion and outlook	309
General discussion		313
7.1	Background	314
7.2	Summary of results	315

7.3	General discussion	321
7.4	Future perspectives	325
7.5	Conclusion	326
Appendices		329
References		348

LIST OF FIGURES

Figure 1.1	Susceptibility loci for the disease phenotypes Crohn's disease (CD) or ulcerative colitis (UC)	6
Figure 1.2	Structure of the colonic mucosa	7
Figure 1.3	Structure of tight junctions and adherens junctions	9
Figure 1.4	Overview of regulatory pathways involved in the cell-mediated immune response and the characteristic defects in the disease phenotypes Crohn's disease (CD) and ulcerative colitis (UC)	12
Figure 1.5	From genes to metabolites in biological systems and the influence of microbial metabolism	20
Figure 1.6	Metabolism of omega-6 and omega-3 polyunsaturated fatty acids (n-6 and n-3 PUFA) from precursor fatty acids	28
Figure 1.7	Intestinal digestion and absorption of dietary lipids	30
Figure 1.8	Putative mechanism of action of omega-3 polyunsaturated fatty acids (n-3 PUFA) on immune cell functions	32
Figure 1.9	Overview of chapters in this dissertation.	39
Figure 2.1	Design for the "EPA time-course experiment"	44
Figure 2.2	Design for the "salmon diet experiment"	49
Figure 2.3	Transcriptomic analysis workflow	58
Figure 2.4	Principle of 2-dimensional (2D) gel electrophoresis	63
Figure 2.5	Proteomic analysis workflow	64
Figure 2.6	Metabolomic analysis workflow	72
Figure 2.7	Microbiomic analysis workflow	78
Figure 3.1	Design for the "EPA time-course experiment"	92
Figure 3.2	Mean body weights (g) for C57BL/6J and <i>Il10</i> ^{-/-} mice over the experimental period	94
Figure 3.3	Histological injury scores (HIS) obtained from the colon of <i>Il10</i> ^{-/-} mice at 9 and 12 weeks of age	97
Figure 3.4	Heatmap of colon gene expression profiles	100
Figure 3.5	Heatmap showing the set of genes comprising the KEGG pathway <i>Tryptophan metabolism</i>	109
Figure 3.6	Metabolism of tryptophan to xanthurenic acid (<i>via</i> kynurenine), tryptamine and serotonin	110
Figure 3.7	Colon gene expression changes in response to the eicosapentaenoic acid (EPA) diet (vs. oleic acid (OA) diet) in <i>Il10</i> ^{-/-} mice at early stages of colitis	113
Figure 3.8	Colon gene expression changes in response to the eicosapentaenoic acid (EPA) diet (vs. oleic acid (OA) diet) in <i>Il10</i> ^{-/-} mice with established colitis	114
Figure 3.9	Heatmap of peripheral blood mononuclear cell (PBMCs) gene expression profiles	117
Figure 3.10	15 highest p-value-ranked canonical pathways in peripheral blood mononuclear cells (PBMCs) from <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice both fed the oleic acid (OA) diet	123
Figure 3.11	Ingenuity pathway for <i>Primary immunodeficiency signalling</i>	124
Figure 3.12	PBMC gene expression changes in response to the eicosapentaenoic acid (EPA) diet (vs. oleic acid (OA) diet) in <i>Il10</i> ^{-/-} mice at early stages of colitis	128
Figure 3.13	PBMC gene expression changes in response to the eicosapentaenoic acid (EPA) diet (vs. oleic acid (OA) diet) in <i>Il10</i> ^{-/-} mice with established colitis	129
Figure 3.14	Colon gene expression associated with early stages of colitis in <i>Il10</i> ^{-/-} mice	131
Figure 3.15	Peripheral blood mononuclear cell (PBMC) gene expression associated with early stages of colitis in <i>Il10</i> ^{-/-} mice	132
Figure 3.16	Colon gene expression associated with established colitis in <i>Il10</i> ^{-/-} mice	133
Figure 3.17	Peripheral blood mononuclear cell (PBMC) gene expression associated with established colitis in <i>Il10</i> ^{-/-} mice	134
Figure 3.18	Partial Least Squares-Discriminant Analysis (PLS-DA) of metabolite fingerprint from urine in positive and negative ionisation mode at 7.1 (T1), 9 (T2), 10.1 (T3) and 12 (T4) weeks of age	138
Figure 3.19	Venn diagram indicating numbers of significantly different ionisation products in the urine of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice	140
Figure 3.20	Venn diagram indicating numbers of significantly different ionisation products in the	

	urine of mice fed the eicosapentaenoic acid (EPA) diet compared to those fed the oleic acid (OA) diet	141
Figure 3.21	Partial Least Squares-Discriminant Analysis (PLS-DA) of metabolite fingerprint from urine of <i>Il10</i> ^{-/-} mice and C57BL/6J mice in positive and negative ionisation mode	142
Figure 3.22	Partial Least Squares-Discriminant Analysis (PLS-DA) of metabolite fingerprint from urine in positive and negative ionisation mode	143
Figure 3.23	Urinary metabolites associated with metabolism of tryptophan <i>via</i> kynurenine and indoleacetaldehyde	148
Figure 3.24	Urinary metabolites associated with metabolism of tryptophan <i>via</i> serotonin	149
Figure 4.1	Design for the “salmon diet experiment”	178
Figure 4.2	Mean body weights (g) for <i>Il10</i> ^{-/-} and C57BL/6J mice over the experimental period	183
Figure 4.3	Representative images of colon sections stained with haematoxylin and eosin	185
Figure 4.4	Scores of individual histological features in colon tissue of <i>Il10</i> ^{-/-} mice fed the 30% salmon diet and 30% control diet	186
Figure 4.5	KEGG pathway gene sets affected in the colon of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice (both fed the AIN-76A diet)	191
Figure 4.6	KEGG pathway gene sets affected in the colon of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice (both fed the 15% control diet)	196
Figure 4.7	Biological interaction network of differentially expressed genes associated with the ten most significant biological functions in <i>Il10</i> ^{-/-} mice fed the 15% salmon diet relative to those fed the 15% control diet	199
Figure 4.8	KEGG pathway gene sets affected in the colon of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice (both fed the 30% control diet)	203
Figure 4.9	Biological interaction network of differentially expressed genes associated with the ten most significant biological functions in the colon of <i>Il10</i> ^{-/-} mice fed the 30% salmon diet compared to those fed the 30% control diet	209
Figure 4.10	Biological interaction network of transcription factors	210
Figure 4.11	KEGG pathway gene sets affected in the colon of <i>Il10</i> ^{-/-} mice fed the 30% salmon diet compared to those fed the 30% control diet	211
Figure 4.12	Ingenuity pathway for <i>Production of Nitric Oxide and Reactive Oxygen Species in macrophages</i>	215
Figure 4.13	KEGG pathway gene sets affected in the colon of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice (both fed the 45% control diet)	216
Figure 4.14	Biological interaction network of differentially expressed genes associated with the ten most significant biological functions in the colon of C57BL/6J mice fed the 45% salmon diet	219
Figure 5.1	Representative gel image with selected spot-features corresponding to proteins for identification	235
Figure 5.2	Ingenuity pathway for <i>Antigen presentation</i> indicating molecular events that lead to the presentation of antigens to CD4 ⁺ and CD8 ⁺ T cells during colitis	242
Figure 5.3	Ingenuity pathway for <i>Antigen presentation</i> indicating molecular events that lead to the presentation of antigens to CD4 ⁺ and CD8 ⁺ T cells in <i>Il10</i> ^{-/-} mice fed the 30% salmon diet	245
Figure 5.4	Biological interaction network of differentially expressed genes associated with the ten most significant biological functions in the liver of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice both fed the 30% control diet	249
Figure 5.5	KEGG pathway gene sets differentially expressed in the liver of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice fed the 30% control diet	250
Figure 5.6	KEGG pathway gene sets differentially expressed in the liver of <i>Il10</i> ^{-/-} mice fed the 30% salmon diet compared to those fed the 30% control diet	253
Figure 5.7	Structural elucidation of xanthurenic acid	257
Figure 5.8	Structural elucidation of xanthurenic acid glucoside	257
Figure 5.9	Structural elucidation of γ -CEHC glucoside	258
Figure 5.10	Structural elucidation of α -CEHC glucuronide	258
Figure 5.11	Partial Least Squares-Discriminant Analysis (PLS-DA) of mouse urine in negative and positive ionisation mode at 6.2, 9 and 11.5 weeks of age	261
Figure 5.12	Peak intensities of the molecular ions corresponding to the urinary metabolites xanthurenic acid glucoside (negative ionisation mode) and xanthurenic acid (positive and negative)	264
Figure 5.13	Chao1 index [325] of caecal microbiota	268

Figure 5.14	Dominant phyla in the caeca of C57BL/6J mice and <i>Il10</i> ^{-/-} mice fed the 30% control diet and the 30% salmon diet	269
Figure 5.15	Principal Coordinate Analysis (PCoA) using (A) an unweighted UniFrac method and (B) a weighted UniFrac method	274
Figure 6.1	Biological pathway for <i>Tryptophan metabolism</i> indicating molecular events that lead to the biosynthesis of melatonin from tryptophan (<i>via</i> serotonin), or xanthurenic acid (<i>via</i> kynurenine) in <i>Il10</i> ^{-/-} mice fed the 30% control diet compared to C57BL/6J mice fed the same diet.	296
Figure 6.2	Correlation of urinary α -CEHC glucuronide (“M453.1T348” and “M472.1T347”) and γ -CEHC glucoside (“M425.1T334” and “M444.1T334”) abundance with hepatic expression of genes	298
Figure 6.3	Relevance network indicating correlation of (A) negative and (B) positive ions and the caecal microbiota, with expression values of <i>Il10</i> ^{-/-} mice fed a 30% control diet (vs. C57BL/6J mice) overlaid	299
Figure 6.4	Biological pathway for <i>Tryptophan metabolism</i> indicating molecular events that lead to the biosynthesis of melatonin from tryptophan (<i>via</i> serotonin), or xanthurenic acid (<i>via</i> kynurenine) in <i>Il10</i> ^{-/-} mice fed the 30% salmon diet compared to those fed the 30% control diet.	301
Figure 6.5	Correlation network of urinary α -CEHC glucuronide (“M453.1T348” and “M472.1T347”) and γ -CEHC glucoside (“M425.1T334” and “M444.1T334”) abundance with hepatic expression of genes	303
Figure 6.6	Relevance network indicating correlation of (A) negative and (B) positive ions and the caecal microbiota, with expression values of <i>Il10</i> ^{-/-} mice fed the 30% salmon diet compared to those fed the 30% control diet overlaid	304
Figure 6.7	Proposed hepatic molecular mechanisms that lead to the elimination of α -tocopherol in the form of α -CEHC glucuronide	311

LIST OF TABLES

Table 1.1	Overview of factors that contribute to the pathogenesis of inflammatory bowel diseases (IBD) and the disease phenotypes ulcerative colitis (UC) and Crohn's disease (CD)	3
Table 1.2	Characteristic changes in microbial community composition in inflammatory bowel diseases (IBD)	13
Table 1.3	Most commonly used mouse models of inflammatory bowel diseases (IBD)	16
Table 1.4	Nutritional profile of farmed New Zealand Chinook salmon fillets (<i>Oncorhynchus tshawytscha</i>).	25
Table 2.1	Formulation of the unmodified AIN-76A diet and AIN-76A-based eicosapentaenoic acid (EPA) and oleic acid (OA) diets for the "EPA time-course experiment"	47
Table 2.2	Formulation of salmon and control diets for the "salmon diet experiment"	52
Table 2.3	Preparation of solutions and buffers used for colon protein analysis	66
Table 2.4	Numbers of urine samples obtained from C57BL/6J mice and <i>Il10</i> ^{-/-} mice fed AIN-76A diets, either unmodified, or enriched with oleic acid (OA) or eicosapentaenoic acid (EPA)	75
Table 2.5	Numbers of urine samples obtained from C57BL/6J mice and <i>Il10</i> ^{-/-} mice fed 30% salmon or 30% control diets	76
Table 2.6	Overview of samples collected and measurements performed in the "EPA time-course experiment" and "salmon diet experiment"	84
Table 2.7	Tasks performed in the "EPA time-course experiment" and "salmon diet experiment" including contributions by the PhD candidate	85
Table 3.1	Analysis of growth performance for "EPA time-course experiment"	93
Table 3.2	Estimated mean histological injury scores (HIS) obtained from intestinal sections of <i>Il10</i> ^{-/-} mice fed AIN-76A diets, either unmodified, or enriched with 3.7% oleic acid (OA) or 3.7% eicosapentaenoic acid (EPA)	96
Table 3.3	Numbers of differentially expressed genes in the colon of <i>Il10</i> ^{-/-} mice and C57BL/6J mice fed AIN-76A diets, either unmodified, or enriched with oleic acid (OA) or eicosapentaenoic acid (EPA)	99
Table 3.4	Genes with the highest fold-changes (FC) between the colon from <i>Il10</i> ^{-/-} mice and C57BL/6J mice fed the oleic acid (OA) diet (9 and 12 weeks of age)	102
Table 3.5	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice fed the oleic acid (OA) diet (9 and 12 weeks of age)	103
Table 3.6	KEGG pathway gene sets affected in the colon of <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice fed the oleic acid (OA) diet and sampled at 9 or 12 weeks of age	106
Table 3.7	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice fed the eicosapentaenoic acid (EPA) diet compared to those fed the oleic acid (OA) diet (9 and 12 weeks of age)	112
Table 3.8	Numbers of differentially expressed genes in peripheral blood mononuclear cells (PBMCs) of <i>Il10</i> ^{-/-} mice and C57BL/6J mice fed the oleic acid (OA) or the eicosapentaenoic acid (EPA) diets	116
Table 3.9	Differentially expressed genes in peripheral blood mononuclear cells (PBMCs) from <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice fed the oleic acid (OA) diet that were in common at 9 and 12 weeks of age	120
Table 3.10	Most significantly affected biological functions in peripheral blood mononuclear cells (PBMCs) from <i>Il10</i> ^{-/-} mice relative to C57BL/6J mice fed the oleic acid (OA) diet (9 and 12 weeks of age)	122
Table 3.11	Differentially expressed genes in peripheral blood mononuclear cells (PBMCs) from <i>Il10</i> ^{-/-} mice	126
Table 3.12	Most significantly affected biological functions in peripheral blood mononuclear cells (PBMCs) from <i>Il10</i> ^{-/-} mice fed the eicosapentaenoic acid (EPA) diet	127
Table 3.13	Summary of gene expression changes in the colon and peripheral blood mononuclear cells (PBMCs) from <i>Il10</i> ^{-/-} and C57BL/6J mice	135
Table 3.14	Permutation MANOVA of the urinary metabolite fingerprints	137
Table 3.15	Positive ionisation products significantly different in the urine from <i>Il10</i> ^{-/-} mice compared to C57BL/6J mice	144
Table 3.16	Negative ionisation products significantly different in the urine from <i>Il10</i> ^{-/-} mice	

	compared to C57BL/6J mice	146
Table 3.17	Positive ionisation products with differential abundance in the urine of mice fed the eicosapentaenoic acid (EPA) diet	151
Table 3.18	Negative ionisation products with differential abundance in the urine of mice fed the eicosapentaenoic acid (EPA) diet	154
Table 3.19	Summary of urinary metabolites detected in <i>Il10</i> ^{-/-} and C57BL/6J mice fed AIN-76A-based diets, either enriched with eicosapentaenoic acid (EPA) or oleic acid (OA)	158
Table 4.1	Nutritional composition of diets salmon and control diets	180
Table 4.2	Analysis of growth performance for “salmon diet experiment”	182
Table 4.3	Numbers of differentially expressed genes in the colon of C57BL/6J and <i>Il10</i> ^{-/-} mice	188
Table 4.4	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice relative to C57BL/6J mice (both fed the AIN-76A diet)	190
Table 4.5	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice relative to C57BL/6J mice (both fed the 15% control diet)	195
Table 4.6	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice fed the 15% salmon diet relative to those fed the 15% control diet	198
Table 4.7	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice relative to C57BL/6J mice (both fed the 30% control diet)	202
Table 4.8	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice fed the 30% salmon diet relative to those fed the 30% control diet	208
Table 4.9	Most significantly affected biological functions in the colon of <i>Il10</i> ^{-/-} mice relative to C57BL/6J mice (both fed the 45% control diet)	213
Table 4.10	Most significantly affected biological functions in the colon of C57BL/6J mice fed the 45% salmon diet relative to those fed the 45% control diet	218
Table 5.1	List of differentially expressed proteins in the colon of <i>Il10</i> ^{-/-} and C57BL/6J mice fed the 30% control or 30% salmon diet	236
Table 5.2	Numbers of differentially expressed genes in the liver of C57BL/6J and <i>Il10</i> ^{-/-} mice	246
Table 5.3	Most significantly affected biological functions in the liver of <i>Il10</i> ^{-/-} relative to C57BL/6J mice (both fed the 30% control diet)	248
Table 5.4	Differentially expressed genes associated with <i>Xenobiotic metabolism</i> in the liver of <i>Il10</i> ^{-/-} mice	251
Table 5.5	Differentially expressed genes associated with <i>Lipid metabolism</i> and <i>Xenobiotics metabolism</i> in the liver of C57BL/6J mice or <i>Il10</i> ^{-/-} mice fed the 30% salmon diet compared to those fed the 30% control diet	254
Table 5.6	Permutation MANOVA of the urinary metabolite fingerprints	260
Table 5.7	Ions with the ten highest Partial Least Squares-Discriminant Analysis (PLS-DA) loadings in <i>Il10</i> ^{-/-} and C57BL/6J mice fed 30% control or 30% salmon diets	262
Table 5.8	Average proportions of dominant genera in caecal digesta	271
Table 7.1	Summary of the main biological functions and microbiota affected in <i>Il10</i> ^{-/-} mice fed diets containing either 3.7% eicosapentaenoic acid (EPA) (“EPA time-course experiment”) or 30% salmon (“salmon diet experiment”)	320

LIST OF APPENDICES

Appendix I	Analysis of lyophilised salmon fillets	329
Appendix II	R codes applied for pre-processing of metabolomics data	330
Appendix III	Histological injury scores (HIS) obtained from the duodenum of C57BL/6J and <i>Il10</i> ^{-/-} mice at 9 and 12 weeks of age	331
Appendix IV	Differentially expressed genes in the colon of <i>Il10</i> ^{-/-} mice fed the oleic acid (OA) diet compared to those fed the AIN-76A diet at 12 weeks of age	332
Appendix V	Molecule shapes and relationship types used by Ingenuity Pathway Analysis	333
Appendix VI	Positive ionisation products with differential abundance in the urine of mice fed the eicosapentaenoic acid (EPA) diet compared to those fed the oleic acid (OA) diet	334
Appendix VII	Negative ionisation products with differential abundance in the urine of mice fed the eicosapentaenoic acid (EPA) diet compared to those fed the oleic acid (OA) diet	336
Appendix VIII	Biological pathway for <i>Tryptophan metabolism</i> indicating molecular events that lead to the biosynthesis of melatonin from tryptophan (<i>via</i> serotonin), or xanthurenic acid (<i>via</i> kynurenine) in <i>Il10</i> ^{-/-} mice fed the oleic acid (OA) diet compared to C57BL/6J mice fed the same diet at 12 weeks of age.	338
Appendix IX	Biological pathway for <i>Tryptophan metabolism</i> indicating molecular events that lead to the biosynthesis of melatonin from tryptophan (<i>via</i> serotonin), or xanthurenic acid (<i>via</i> kynurenine) in <i>Il10</i> ^{-/-} mice fed the eicosapentaenoic acid (EPA) diet compared to those fed the oleic acid (OA) diet.	340
Appendix X	Histology scores from (A) C57BL/6J mice and (B) <i>Il10</i> ^{-/-} mice fed diets supplemented with 15%, 30% or 45% salmon and corresponding macronutrient-matched control diets	342
Appendix XI	qPCR validation of microarray results	343
Appendix XII	Unidentified ions with the ten highest Partial Least Squares-Discriminant Analysis (PLS-DA) loadings (“discriminant ions”) in <i>Il10</i> ^{-/-} and C57BL/6J mice fed 30% control or 30% salmon diets	345
Appendix XIII	Proteins identified from previous experiments.	347

LIST OF ABBREVIATIONS

AA	Arachidonic acid
ALA	Alpha-linolenic acid
AMP	Antimicrobial protein
ANOVA	Analysis of variance
CAM	Cell adhesion molecule
CD	Crohn's disease
CEBPB, CEBPD, CEBPE	CCAAT/enhancer binding protein (alpha, beta, delta)
CEHC	Carboxyethylhydroxychroman
CFU	Colony-forming units
CIF	Complex intestinal microbiota
DHA	Docosahexaenoic acid
DIGE	Difference gel electrophoresis
DPA	Docosapentaenoic acid
DSS	Dextran Sodium Sulfate
EF	<i>Enterococcus faecalis</i> and <i>E. faecium</i>
EF×CIF	Solution for bacterial inoculation
EPA	Eicosapentaenoic acid
ESI	Electrospray ionisation
FC	Fold-change
FDR	False discovery rate
GIT	Gastrointestinal tract
GLA	Gamma-linolenic acid
GSEA	Gene Set Enrichment Analysis
GWAS	Genome-Wide Association Study
HE	Haematoxylin and eosin
HIS	Histological injury score
HMDB	Human Metabolome Database
IBD	Inflammatory bowel diseases
IEF	Isoelectric focussing
IL[number]	Interleukin [number]
<i>Il10^{-/-}</i>	Interleukin-10 gene-deficient
IPA	Ingenuity Pathway Analysis
KEGG	Kyoto Encyclopedia of Genes and Genomes
LA	Linoleic acid
LC	Long-chain
LPS	Lipopolysaccharide
LysoPC	Lysophosphatidylcholine

LysoPE	Lysophosphatidylethanolamine
<i>m/z</i>	Mass-per-charge ratio
MANOVA	Multivariate ANOVA
MDT	Marine-derived tocopherol
MS	Mass spectrometry
n-3 PUFA	Omega-3 polyunsaturated fatty acid
n-6 PUFA	Omega-6 polyunsaturated fatty acid
NCBI	National Center for Biotechnology Information
NKT cell	Natural killer T cell
NMR	Nuclear magnetic resonance
OA	Oleic acid
OTU	Operational taxonomic unit
PAMP	Pathogen-associated molecular pattern
PBMC	Peripheral blood mononuclear cell
PBS	Phosphate-buffered saline
PC	Principal coordinate
PCA	Principal Component Analysis
PCoA	Principal Coordinate Analysis
pI	Isoelectric point
PLS-DA	Partial Least Squares-Discriminant Analysis
PPAR	Peroxisome proliferator-activated receptor
PRR	Pattern recognition receptor
QIIME	Quantitative Insights Into Microbial Ecology
qRT-PCR	Real-time reverse transcription polymerase chain reaction
REML	Restricted maximum likelihood
REST	Relative Expression Software Tool
RIN	RNA integrity number
ROS	Reactive Oxygen Species
RT	Retention time
SCFA	Short-chain fatty acid
SDS	Sodium dodecyl sulphate
SDS-PAGE	Sodium dodecyl sulphate polyacrylamide gel electrophoresis
SNP	Single nucleotide polymorphisms
TG	Triacylglycerol
Th cell	T helper cell
TMAO	Trimethylamine N-oxide
TNBS	Trinitrobenzenesulfonic acid
Treg	Regulatory T helper cell
UC	Ulcerative colitis